

Iron and Iron Deficiency

Iron and why we need it

Iron is a mineral needed by our bodies. Iron is a part of all cells and does many things in our bodies. For example, iron (as part of the protein hemoglobin) carries oxygen from our lungs throughout our bodies. Having too little hemoglobin is called anemia. Iron also helps our muscles store and use oxygen.

Iron is a part of many enzymes and is used in many cell functions. Enzymes help our bodies digest foods and also help with many other important reactions that occur within our bodies. When our bodies don't have enough iron, many parts of our bodies are affected.

Iron deficiency and why it is a concern

Iron deficiency is a condition resulting from too little iron in the body. Iron deficiency is the most common nutritional deficiency and the leading cause of anemia.¹

The terms anemia, iron deficiency, and iron deficiency anemia often are used interchangeably but equivalent. Iron deficiency ranges from depleted iron stores without functional or health impairment to iron deficiency with anemia, which affects the functioning of several organ systems.²

Iron deficiency is a concern because it can:

- Iron deficiency can delay normal infant motor function (normal activity and movement) or mental function (normal thinking and processing skills).³⁻⁶
- Iron deficiency anemia during pregnancy can increase risk for small or early (preterm) babies.⁷⁻⁸ Small or early babies are more likely to have health problems or die in the first year of life than infants who are born full term and are not small.
- Iron deficiency can cause fatigue that impairs the ability to do physical work in adults.⁹⁻¹⁰ Iron deficiency may also affect memory or other mental function in teens.¹¹

Causes Of Iron Deficiency

Iron deficiency has many causes. These causes fall into two main categories:

1. Increased iron needs

Many common conditions can cause people to need additional iron:

- Because of their rapid growth, infants and toddlers need more iron than older children. Sometimes it can be hard for them to get enough iron from their normal diet.
- Women who are pregnant have higher iron needs. To get enough, most women must take an iron supplement as recommended by their healthcare provider.
- When people lose blood, they also lose iron. They need extra iron to replace what they have lost. Increased blood loss can occur with heavy menstrual periods, frequent blood donation, as well as with some stomach and intestinal conditions (food sensitivity, hookworms.)

2. Decreased iron intake or absorption (not enough iron taken into the body)

The amount of iron absorbed from the diet depends on many factors:

- Iron from meat, poultry, and fish (i.e., heme iron) is absorbed two to three times more efficiently than iron from plants (i.e., non-heme iron).
- The amount of iron absorbed from plant foods (non-heme iron) depends on the other types of foods eaten at the same meal.
- Foods containing heme iron (meat, poultry, and fish) enhance iron absorption from foods that contain non-heme iron (e.g., fortified cereals, some beans, and spinach).
- Foods containing vitamin C (see Dietary Sources of vitamin C) also enhance non-heme iron absorption when eaten at the same meal.
- Substances (such as polyphenols, phytates, or calcium) that are part of some foods or drinks such as tea, coffee, whole grains, legumes and milk or dairy products can decrease the amount of non-heme iron absorbed at a meal. Calcium can also decrease the amount heme-iron absorbed at a meal. However, for healthy individuals who consume a varied diet that conforms to the Dietary Guidelines for Americans, the amount of iron inhibition from these substances is usually not of concern.
- Vegetarian diets are low in heme iron, but careful meal planning can help increase the amount of iron absorbed.
- Some other factors (such as taking antacids beyond the recommended dose or medicine used to treat peptic ulcer disease and acid reflux) can reduce the amount of acid in the stomach and the iron absorbed and cause iron deficiency.

Increased Iron Needs	Decreased Iron Intake and Absorption
<ul style="list-style-type: none"> • Rapid growth • Pregnancy • Blood loss <ul style="list-style-type: none"> ○ Heavy menstrual periods ○ Frequent blood donation ○ Some stomach and intestinal conditions (food sensitivity, hookworms) 	<ul style="list-style-type: none"> • Lack of heme iron sources in the diet (e.g., vegetarian diets) • Low absorption <ul style="list-style-type: none"> ○ Taking antacids beyond the recommended dose or medicine used to treat peptic ulcer disease and acid reflux can reduce the amount of iron absorbed in the stomach.

People at risk

- Young children and pregnant women are at higher risk of iron deficiency because of rapid growth and higher iron needs.
- Adolescent girls and women of childbearing age are at risk due to menstruation.
- Among children, iron deficiency is seen most often between six months and three years of age due to rapid growth and inadequate intake of dietary iron. Infants and children at highest risk are the following groups:
 - Babies who were born early or small.
 - Babies given cow's milk before age 12 months.
 - Breastfed babies who after age 6 months are not being given plain, iron-fortified cereals or another good source of iron from other foods.
 - Formula-fed babies who do not get iron-fortified formulas.

- Children aged 1–5 years who get more than 24 ounces of cow, goat, or soymilk per day. Excess milk intake can decrease your child's desire for food items with greater iron content, such as meat or iron fortified cereal.
- Children who have special health needs, for example, children with chronic infections or restricted diets.

Signs and Symptoms of Iron Deficiency

Too little iron can impair body functions, but most physical signs and symptoms do not show up unless iron deficiency anemia occurs. Someone with early stages of iron deficiency may have no signs or symptoms. This is why it is important to screen for too little iron among high risk groups.

Signs of iron deficiency anemia include¹²

- Feeling tired and weak
- Decreased work and school performance
- Slow cognitive and social development during childhood
- Difficulty maintaining body temperature
- Decreased immune function, which increases susceptibility to infection
- Glossitis (an inflamed tongue)

Detection of iron deficiency detected

Your doctor or healthcare provider will do blood tests to screen for iron deficiency. No single test is used to diagnose iron deficiency. The most common tests for screening are

- Hemoglobin test (a test that measures hemoglobin which is a protein in the blood that carries oxygen)
 - Hematocrit test (the percentage of red blood cells in your blood by volume)
- These tests show how much iron is in your body. Hemoglobin and hematocrit levels usually aren't decreased until the later stages of iron deficiency, i.e., anemia.

Sometimes other blood tests are used to confirm that anemia is due to iron deficiency. These might include

- Complete blood count (to look at the number and volume of the red blood cells)

- Serum ferritin (a measure of a stored form of iron)
- Serum iron (a measure of the iron in your blood)
- Transferrin saturation (a measure of the transported form of iron)
- Transferrin receptor (a measure of increased red blood cell production)

Treatment of iron deficiency

- If you are found to have an iron deficiency, it is important to see your healthcare provider for treatment. Your treatment will depend on factors such as your age, health, and cause of iron deficiency.
- If your doctor or health care provider thinks that you have iron deficiency she or he may prescribe iron supplements for you to take and then ask that you return after a period to have your hemoglobin or hematocrit tested.
- If your healthcare provider determines that the iron deficiency is due to a diet low in iron, you might be told to eat more iron-rich foods. Your health care provider may also prescribe an iron supplement for you.

Again, it is important to be diagnosed by your healthcare provider because iron deficiency can have causes that aren't related to your diet. Your healthcare provider's recommendations will be specific to your needs.

Prevention of iron deficiency

In general, you can eat a healthful diet that includes good sources of iron. A healthful diet includes fruits, vegetables, whole grains, fat free or nonfat milk and milk products, lean meats, fish, dry beans, eggs, nuts, and is low in saturated fat, trans fats, cholesterol, salt, and added sugars.

In addition to a healthful diet that includes good sources of iron, you can also eat foods that help your body absorb iron better. For example, you can eat a fruit or vegetable that is a good source of vitamin C with a food or meal that contains non-heme iron. Vitamin C helps your body absorb the non-heme iron foods you eat, especially when the food containing non-heme iron and the vitamin-C rich food are eaten at the same meal.

The following recommendations are for specific groups who are at greater risk for iron deficiency.

Babies

- If possible, breastfeed your baby for at least 12 months and starting at 4 to 6 months of age, give your baby plain, iron-fortified infant cereal and/or pureed meat. Just two or more servings a day can meet a baby's iron needs at this age. Meats should be home prepared or commercially prepared plain pureed (chopped until smooth in a blender) meats.
- When your baby is about 6 months of age, include a feeding per day of foods rich in vitamin C with foods that are rich in non-heme iron to improve iron absorption.
- If you can't breastfeed, use iron-fortified formula.
- Don't give low-iron milks (e.g. cow's milk, goat's milk, and soy milk) until your baby is at least 12 months old.
- If your baby was born early or small, talk to your doctor about giving iron drops to your baby.
- If your baby can't get two or more servings per day of iron rich foods (such as iron-fortified cereal or pureed meats), talk to your doctor about giving iron drops to your baby.

Young children (aged 1–5 years)

- After your child is one year old, give no more than three 8 ounce servings of whole cow, goat, or soy milk per day. After your child is 2 years old, low fat or nonfat milks should be used in place of whole milks. Vitamin D-fortified milk is a good source of calcium and vitamin D, but not iron.
- Give your child a diet with iron-rich foods such as iron-fortified breads and iron-fortified cereals and lean meats.
- Include fruits, vegetables or juices that are rich in vitamin C. Vitamin C helps your child absorb non-heme iron especially when the food that is a source of non-heme iron and the vitamin C-rich food are eaten at the same meal.

Adolescent girls and women of childbearing age

- Eat iron-rich foods..
- Eat foods that are vitamin C sources. Vitamin C helps your body absorb non-heme iron especially when the food that is a source of non-heme iron and the vitamin C-rich food are eaten at the same meal.
- Eat lean red meats, poultry, and fish. The iron in these foods is easier for your body to absorb than the iron in plant foods.

Pregnant women

- Eat iron-rich foods.
- Eat foods that are vitamin C sources. Vitamin C helps your body absorb non-heme iron especially when the food that is a source of non-heme iron and the vitamin-C rich food are eaten at the same meal.
- Eat lean red meats, poultry, and fish. The iron in these foods is easier for your body to absorb than the iron in plant foods.
- Talk to your doctor about taking an iron supplement.

Recommended Dietary Allowance (RDA) for iron

If you have already been diagnosed with iron deficiency, talk to your doctor or healthcare provider about treatment. For healthy individuals, the Recommended Dietary Allowance (RDA) for iron is listed in the following table.

**RECOMMENDED DIETARY ALLOWANCES FOR INDIANS
(Macronutrients and Minerals)**

Group	Particulars	Body wt. kg	Net Energy Kcal/d	Protein g/d	Visible Fat g/day	Calcium mg/d	Iron mg/d
Man	Sedentary work	60	2320	60	25	600	17
	Moderate work		2730		30		
	Heavy work		3490		40		
Woman	Sedentary work	55	1900	55	20	600	21
	Moderate work		2230		25		
	Heavy work		2850		30		
	Pregnant woman		+350	82.2	30	1200	3 5
	Lactation 0-6 months		+600	77.9	30	1200	25
	6-12 months		+520	70.2	30		
Infants	0-6 months	5.4	92 Kcal/kg/d	1.16 g/kg/d	–	500	--
	6-12 months	8.4	80 Kcal/kg/d	1.69 g/kg/f	19		46 µg/kg/day
Children	1-3 years	12.9	1060	16.7	27	600	09
	4-6 years	18	1350	20.1	25		13
	7-9 years	25.1	1690	29.5	30		16
Boys	10-12 years	34.3	2190	39.9	35	800	21
Girls	10-12 years	35.0	2010	40.4	35	800	27
Boys	13-15 years	47.6	2750	54.3	45	800	32
Girls	13-15 years	46.6	2330	51.9	40	800	27
Boys	16-17 years	55.4	3020	61.5	50	800	28
Girls	16-17 years	52.1	2440	55.5	35	800	26

References

- Centers for Disease Control and Prevention. Iron deficiency – United States, 1999–2000. *MMWR* 2002;51:897–899.
- Akman M, Cebeci D, Okur V, Angin H, Abali O, Akman AC. The effects of iron deficiency on infants' developmental test performance. *Acta Paediatr.* 2004 Oct;93(10):1391–6.
- Friel JK, Aziz K, Andrews WL, Harding SV, Courage ML, Adams RJ. A double-masked, randomized control trial of iron supplementation in early infancy in healthy term breast-fed infants. *J Pediatr.* 2003 Nov;143(5):582–6.
- Lozoff B, De Andraca I, Castillo M, Smith JB, Walter T, Pino P. Behavioral and developmental effects of preventing iron-deficiency anemia in healthy full-term infants. *Pediatrics.* 2003 Oct;112(4):846–54.

5. Grantham-McGregor S, Ani C. A review of studies on the effect of iron deficiency on cognitive development in children. *J Nutr.* 2001 Feb;131(2S-2):649S-666S; discussion 666S-668S.
6. Ronnenberg AG, Wood RJ, Wang X, Xing H, Chen C, Chen D, Guang W, Huang A, Wang L, Xu X. Preconception hemoglobin and ferritin concentrations are associated with pregnancy outcome in a prospective cohort of Chinese women. *J Nutr.* 2004 Oct;134(10):2586-91.
7. Scholl TO, Hediger ML, Fischer RL, Shearer JW. Anemia vs iron deficiency: increased risk of preterm delivery in a prospective study. *Am J Clin Nutr.* 1992 May;55(5):985-8.
8. Brownlie T 4th, Utermohlen V, Hinton PS, Haas JD. Tissue iron deficiency without anemia impairs adaptation in endurance capacity after aerobic training in previously untrained women. *Am J Clin Nutr.* 2004 Mar;79(3):437-43.
9. Haas JD, Brownlie T 4th. Iron deficiency and reduced work capacity: a critical review of the research to determine a causal relationship. *J Nutr.* 2001 Feb;131(2S-2):676S-688S; discussion 688S-690S.
10. Bruner AB, Joffe A, Duggan AK, Casella JF, Brandt J. Randomised study of cognitive effects of iron supplementation in non-anaemic iron-deficient adolescent girls. *Lancet.* 1996 Oct 12;348(9033):992-6.
11. US National Library of Medicine, NIH. Iron deficiency anemia. http://www.nlm.nih.gov/health/dci/Diseases/ida/ida_what.html
12. Office of Dietary Supplements, NIH. Dietary supplement fact sheet. Available online: <http://dietary-supplements.info.nih.gov/factsheets/iron.asp>
13. Dietary Guidelines for Indians. A manual. National Institute of Nutrition, Hyderabad, A.P.